

## What is special about the vulvar vestibule?

Melissa A. Farmer

**AQ:1** **T**he reproductive machinery is evolutionarily precious. As a result, physiological mechanisms evolved to protect reproductive organs from common environmental threats. Ongoing exposure to potentially pathogenic fungi, bacteria, and viruses has made the female lower reproductive tract a veritable battleground for immune defense. The external vulva and vaginal canal express a unique immune profile that differs from other mucosal sites and elsewhere in the periphery, and this specialization is preserved across species.<sup>10</sup> It therefore follows that pain induced by inflammation may also show vulvovaginal-specific properties.

The elegant work presented by Foster et al.<sup>6</sup> expands our understanding of the complex relationship between pain and immune function with the novel observation that the vulvar vestibule—a thin band of tissue demarcating the entrance to the vagina—exhibits a highly localized and tissue-specific proinflammatory response in healthy women. This response is further amplified in women with localized provoked vulvodynia (LPV), a common type of chronic pelvic pain that afflicts 9% to 16% of premenopausal women.<sup>8</sup> Localized provoked vulvodynia is characterized by sharp, burning pain localized to the vulvar vestibule in response to light pressure (ie, mechanical allodynia) and is accompanied by enhanced vulvar pain perception (ie, mechanical hyperalgesia) and continued pain that is temporally dissociated from its stimulus (ie, “after sensation”). The mechanisms underlying LPV remain unknown, despite equivocal evidence for low-grade vulvar inflammation, altered peptidergic vulvar innervation, and genetic susceptibilities that contribute to abnormal inflammatory cascades.

*Candida albicans* is a known inflammogen across species and is implicated in more than 90% of vulvovaginal yeast infections in women. The high incidence of recurrent yeast infections in women with LPV prompted the hypothesis that yeast-induced vulvovaginal inflammation may underlie peripheral (and potentially central) sensitization of primary afferent nerve endings at the vestibule. This hypothesis allows for experience-dependent inflammatory insults to play a role in the initiation and/or maintenance of LPV (supported by direct activation of nociceptors by pathogenic bacteria<sup>3</sup>), as well as genetic polymorphisms that are frequently identified in this population.<sup>2</sup> The former point is supported by an experimental mouse model of LPV that confirmed that repeated vulvovaginal infections with *C. albicans* are sufficient to cause chronic vulvar pain behavior and increased vulvar peptidergic innervation, a finding that parallels

observations in the clinical literature.<sup>2,4</sup> Furthermore, some women with chronic vulvar pain may be exceptionally vulnerable to *C. albicans* exposure, as indicated by enhanced contact sensitivity to intradermal *C. albicans* injections and increased fibroblast-mediated cytokine production.<sup>7,11</sup> The question how the vestibule uniquely mediates these symptoms remains, and how can we differentiate normal vulvar nociceptive processes from pathological ones?

To address these questions, Foster et al.<sup>6</sup> evaluated women with and without chronic vulvar pain with vulvar quantitative sensory testing and examined vulvar tissue variability in fibroblast-mediated cytokine release. Women were assessed for alterations in pain perception associated with vulvar mechanical allodynia (through algometry) and hyperalgesia (through a cotton swab test). Mechanical pain threshold and suprathreshold levels were measured at the vestibule and an adjacent vulvar control site, and tissue biopsies were obtained from the same regions. In normal vulvar tissue of healthy women, fibroblast exposure to *Candida* and non-*Candida* yeast elicited modest fibroblast-mediated inflammation, as measured by interleukin-6 (IL-6) and prostaglandin E2 (PGE2) levels. Importantly, a robust increase in the magnitude of this proinflammatory response was evident at the vestibule, compared with the adjacent vulvar tissue. These observations strongly support that the vulvar vestibule is an immunologically unique tissue that exhibits enhanced inflammatory responsiveness to multiple genera of fungi, mediated in part by specialized fibroblasts. In women with LPV, exaggerated IL-6 and PGE2 levels were observed at both vulvar sites after *Candida* exposures. Vestibule fibroblasts, in particular, induced a profound increase of proinflammatory mediators in response to all yeast stimuli. Therefore, the vulvar tissues of women with and without LPV were differentiated by the degree of naturally occurring inflammation.

Given the increasing focus on peripheral and central immune factors underlying acute and chronic pain, this work is a timely and topical advance in our understanding of site-specific mucosal immunity. The unique inflammatory properties of the vestibule seem to facilitate rapid detection of pathogens at a critical anatomical gateway to the reproductive tract. It is feasible that other structures derived from the endoderm, including the bladder and anterior urethral wall, may also exhibit unique immune properties. Notably, the high incidence of comorbid vulvar and bladder pain in young women, in some cases, may reflect vulnerabilities of endoderm-derived tissue.<sup>1,5</sup> These possibilities require a shift in our understanding of inflammation because it relates to pain pathology, a sort of immunopathological relativism: the clinical significance of inflammation depends on its anatomical context. For instance, the magnitude of a proinflammatory response—rather than its presence alone—may be critical in differentiating pelvic pain populations from healthy individuals, and from one another. Long after the original nociceptive stimuli are gone, in vivo and in vitro

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

Department of Physiology, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA

PAIN 00 (2015) 1–2

© 2015 International Association for the Study of Pain

<http://dx.doi.org/10.1097.XXX>

**AQ:2** pathogen/irritant challenges that evoke proinflammatory responses can provide an ideal diagnostic strategy for examining such responses.

Clinically, the correlations between self-reported vulvar pain sensitivity and distinct fibroblast-mediated proinflammatory factors reported by Foster et al.<sup>6</sup> provide some tantalizing results. Matching such specialized immune mechanisms to clinical symptoms is the closest the field has come to in identifying an objective marker of pathology of LPV. This finding in part reflects the authors' painstaking efforts to obtain tissue from patient-identified sites of vulvar pain. Furthermore, the data presented in Tables 2 and 3 raise the possibility that levels of IL-6 and PGE2 are associated with distinct sensory aspects of LPV. Fibroblast-mediated IL-6 levels seem to uniquely relate to pain threshold and, by extension, mechanical allodynia, whereas PGE2 production tended to preferentially reflect suprathreshold pain evoked during the cotton swab test (8/10 mean pain rating in this sample). These speculations, as well as other novel hypotheses raised by the authors, would require larger samples of women with LPV who are carefully phenotyped. The likely existence of distinct subtypes of LPV suggests that a principle-components analysis would be needed to identify women who show sensitized inflammatory responses, because any clinical sample may contain mechanistically distinct variants of LPV.<sup>9</sup> Furthermore, the use of zymosan as an inflammatory stimulus can be applied across laboratories without the need for biohazard safety measures required for handling *Candida*. This convenience facilitates its use in animal model development to better understand the acute and chronic impact of vulvovaginal yeast vs bacterial exposure, the genetic mediation of this impact, and prevention and treatment of resulting vulvar pain behavior.

Without mincing words, I believe that the proof of concept of Foster et al. is one of the most significant advances in vulvodynia research over the past 20 years. Their work answers long-standing questions about the significance of vulvar inflammation in women with and without LPV pain, establishes tissue-specific fibroblasts as sufficient and sensitive mediators of cytokine release, and relates fibroblast-mediated cytokine levels with clinically meaningful behavioral outcomes used in the assessment and diagnosis of LPV. This work is the closest we have come to in identifying objective and quantifiable indicators of peripheral pathology underlying persistent vulvar pain. Moving forward, it will be important to integrate these findings with information on peripheral afferent sensitization, spinally mediated central sensitization, and supraspinal processes underlying emotional learning of the chronic pain state.

## Conflict of interest statement

The author has no conflicts of interest to declare.

M. A. Farmer is supported by the National Vulvodynia Association, NICHD L30HD077854, and NIDDK U01 DK082342, in collaboration with the Multidisciplinary Approach to the Study of Chronic Pelvic Pain (MAPP) Research Network.

## Article history:

Received 5 January 2015

Received in revised form 7 January 2015

Accepted 7 January 2015

Available online XXXX

## References

- [1] Bogart LM, Berry SH, Clemens JQ. Symptoms of interstitial cystitis, painful bladder syndrome, and similar diseases in women: a systematic review. *J Urol* 2007;177:450–6.
- [2] Bohm-Starke N. Medical and physical predictors of localized provoked vulvodynia. *Acta Obstet Gynecol Scand* 2010;89:1504–10.
- [3] Chiu IM, Heesters BA, Ghasemlou N, Von Hehn CA, Zhao F, Tran J, Wainger B, Strominger A, Muralidharan S, Horswill AR, Bubeck-Wardenburg J, Hwang SW, Carroll MC, Woolf CJ. Bacteria activate sensory neurons that modulate pain and inflammation. *Nature* 2013;501:52–7.
- [4] Farmer MA, Taylor AM, Bailey AL, Tuttle AH, MacIntyre LC, Milagrosa ZE, Crissman HP, Bennett GJ, Ribeiro-da-Silva A, Binik YM, Mogil JS. Repeated vulvovaginal fungal infections cause persistent pain in a mouse model of vulvodynia. *Sci Transl Med* 2011;3:101ra91.
- [5] Fitzpatrick CC, DeLancey JO, Elkins TE, McGuire EJ. Vulvar vestibulitis and interstitial cystitis: a disorder of urogenital sinus-derived epithelium? *Obstet Gynecol* 81:860–2. **AQ:3**
- [6] Foster CD, Megan L., Falsetta ML, Woeller CF, Pollock SJ, Song K, Bonham A, Haidaris CG, Stodgell CJ, Messing SP, Iadarola M, Phipps RP. Site-specific mesenchymal control of inflammatory pain to yeast challenge in vulvodynia afflicted and pain-free women. *PAIN* 2015. **AQ:4**
- [7] Foster DC, Piekarz KH, Murant TI, LaPoint R, Haidaris CG, Phipps RP. Enhanced synthesis of proinflammatory cytokines by vulvar vestibule fibroblasts: Implications for vulvar vestibulitis. *Am J Obstet Gynecol* 2007; 196:346.e1–8.
- [8] Haefner HK, Collins ME, Davis GD, Edwards L, Foster DC, Hartmann ED, Kaufman RH, Lynch PJ, Margesson LJ, Moyal-Barracco M, Piper CK, Reed BD, Stewart EG, Wilkinson EJ. The vulvodynia guideline. *J Low Genit Tract Dis* 2005;9:40–51.
- [9] Nguyen RH, Veasley C, Smolenski D. Latent class analysis of comorbidity patterns among women with generalized and localized vulvodynia: preliminary findings. *J Pain Res* 2013;6:303–9.
- [10] Quayle AJ. The innate and early immune response to pathogen challenge in the female genital tract and the pivotal role of epithelial cells. *J Reprod Immunol* 2002;57:61–79.
- [11] Ramirez De Knott HM, McCormick TS, Do SO, Goodman W, Ghannoum MA, Cooper KD, Nedorost ST. Cutaneous hypersensitivity to *Candida albicans* in idiopathic vulvodynia. *Contact Dermatitis* 2005;53:214–8.